

	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	---

[illegible]

Description

DI ^{Insert} ~~act~~
The present invention concerns a new growth/ differentiation factor of the TGF- β family and DNA sequences coding therefor.

The BMP-, TGF- and inhibin-related proteins are members of the TGF- β family of growth factors (Roberts and Sporn, Handbook of Experimental Pharmacology 95, 419-472 (1990)). They are relevant for a wide range of medical therapeutic methods and applications. These factors are suitable for methods relating to wound healing and tissue regeneration. Moreover several members of the TGF- β family induce tissue growth for example the growth of bones.

Wozney (Progress in Growth Factor Research 1 (1989), 267-280) and Vale et al. (Handbook of Experimental Pharmacology 95 (1990), 211-248) describe various growth factors for example those which are related to the BMP and the activin/inhibin group. The members of this group have significant structural similarities. The precursor of the protein is composed of an amino-terminal signal sequence, a propeptide sequence and a carboxy-terminal sequence of 110 to 140 amino acids which is cleaved from the precursor and represents the mature protein. Furthermore its members are defined by an amino acid sequence homology. The mature protein contains the sequences that are conserved most, in particular seven cysteine residues which are conserved among the family members. The TGF- β -like proteins are multifunctional, hormonally active growth factors. They also have related biological activities for example chemotactic attraction of cells, promotion of cell differentiation and tissue-inducing capabilities. EP 0 222 491 A1 discloses sequences of inhibin alpha and beta chains.

On the whole the proteins of the TGF- β family show differences in their structure which leads to considerable variations in

A nucleotide and amino acid sequence for the TGF- β protein MP121 is given in the patent application PCT/EP93/00350 in which a major part of the sequence corresponding to the mature protein is stated. The complete sequence of the propeptide MP121 is not disclosed.

The underlying object of the present invention is to provide DNA sequences which code for new members of the TGF- β protein family with mitogenic and/or differentiation-inductive potential. The object of the present invention is in

SEQ ID NO.3 shows the complete nucleotide sequence of the DNA coding for the TGF- β protein MP121 from the mouse. The coding region begins at the ATG start codon at nucleotide 131 and ends at the stop codon beginning at position 1107. The start of the mature protein preferably begins at nucleotide 839. A ca. 5.5 kb large intron is located in the genomic DNA between position 446 and 447.

SEQ ID NO. 4 shows the complete amino acid sequence of the preproprotein of the TGF- β protein MP121 from the mouse which has been derived from the nucleotide sequence shown in SEQ ID NO. 3. The mature protein begins in the region of amino acids 217-240 in analogy to the human MP121 of SEQ ID NO.2. It is most preferred when the mature protein starts at amino acid 237 so that the mature part consists of 116 amino acids as in the human MP121. Members of the TGF- β family are frequently cleaved behind a RXXR cleavage site in order to separate the mature part from the precursor (see Özkaynak et al., J. Biol. Chem. 267, 25220-25227 (1992) and the literature cited therein). In the case of MP121 from the mouse it is also conceivable that the beginning of the mature protein is at least sometimes at amino acid 236.

SEQ ID NO.5 shows the nucleotide sequence of the human MP121 gene at the exon/intron junctions. The nucleotides from both exons are marked by capital letters those of the intron by small letters.

Figure 1 shows a comparison of the amino acid sequence of human MP121 with some members of the TGF- β family (inhibin α and β chains) starting at the first of the seven conserved cysteine residues. * denotes that the amino acid is the same in all compared proteins; + denotes that the amino acid

corresponds in at least one of the proteins compared to human MP121.

Sub D4
Figure 2 shows the nucleotide sequences of the oligonucleotide primers which were used in the present invention and a comparison of these sequences with known members of the TGF- β family. M denotes A or C, S denotes C or G, R denotes A or G and K denotes G or T. 2a shows the sequence of primer OD, 2b shows the sequence of primer OID.

Sub C2
Figure 3 shows a diagram of a Western blot using chicken antibodies against human MP121.

Sub C3
Figure 4 shows the expression of MP121 compared to activin β_A and β_B in various mouse tissues.

Sub C4
Figure 5 shows a positive influence on the survival of dopaminergic neurones by treatment with partially purified MP121.

Sub C5
Figure 6 shows a Western blot using rabbit antibodies against human MP121.

Sub C6
Figure 7 shows the stimulation of nerve fibre outgrowth from the embryonic retina by treatment with partially purified MP121.

Sub C7
Figure 8 shows that partially purified MP121 can inhibit EGF induced DNA synthesis in hepatocytes.

Sub C8
Figure 9 shows the influence of partially purified MP121 on erythroid differentiation.

Within the scope of the present invention the term "mature protein" also encompasses functional partial regions of the complete protein which exhibit essentially the same biological activity and preferably those partial regions which include at

least the region of the seven cysteines that are conserved in the TGF- β family. In this case it is in particular possible that the N-terminus of the mature protein is slightly modified i.e. deviates from the sequences shown in SEQ ID NO.2 and 4. In this connection additional amino acids, which do not influence the functionality of the protein, may be present or amino acids may be absent provided that in this case the functionality is also not impaired. However, it is preferred that the human protein and the mouse protein contain all amino acids starting with amino acid 237 of the amino acid sequence shown in SEQ ID NO.2 and SEQ ID NO.4. It is already known from other family members of the TGF- β family that the attachment of additional amino acids to the N-terminus of the mature protein does not influence the activity wherein inter alia 6 additional histidines were attached to the N-terminus.

Therefore the present invention encompasses the part coding for the mature protein in accordance with the above-mentioned definition and if necessary, further functional parts of the nucleotide sequence shown in SEQ ID NO. 1 as well as sequences that correspond to this sequence within the scope of the degeneracy of the genetic code and allelic derivatives of such sequences. Furthermore the present invention also encompasses DNA sequences which code for a protein of the TGF- β family which were obtained from other mammals and which have a sequence that deviates slightly due to their origin but which, however, code for proteins having in principle the same biological function and also sequences that differ only slightly. Such sequences correspond to one another to a very large extent as can be seen by comparing SEQ ID NO. 1 and NO. 3.

In addition the present invention also covers sequences hybridizing with such sequences provided that such a DNA molecule at least completely contains the part coding for a mature protein of the TGF- β family (according to the above definition) and the biological activity is retained.

00504383-104000

On the other hand the DNA molecules can also include functional signal or/and propeptide parts of other proteins e.g. of proteins with the cystine knot motif (Cell, vol. 73 (1993), p. 421-424) and in particular of other proteins of the TGF- β family e.g. the abovementioned activin/inhibin or BMP proteins especially also MP52 (see PCR/EP94/02630) in addition to the part coding for the mature protein. The respective nucleotide sequences can be found in the aforementioned references to the disclosure of which reference is herewith made. In this case it is important that the correct reading frame for the mature protein is preserved. Depending in which host cells expression takes place, the presence of another

Although the allelic, degenerated and hybridizing sequences and sequences derived from other vertebrates which are covered by the present invention have structural differences due to slight changes in the nucleotide or/and amino acid sequence, proteins which are coded by such sequences still essentially have the same useful properties which enable them to be used in essentially the same medical fields of application.

According to the present invention the term "hybridization" denotes the usual hybridization conditions, preferably conditions with a salt concentration of 6 x SSC at 62 to 66°C followed by a one hour wash with 0.6 x SSC, 0.1 % SDS at 62 to 66°C.

Preferred embodiments of the present invention are DNA sequences as defined above which are obtainable from vertebrates, preferably mammals such as pigs, cows and rodents such as rats or mice and in particular from primates such as humans or which are copied from corresponding sequences.

A particularly preferred embodiment of the present invention are the sequences shown in SEQ ID NO. 1 and 3 and denoted human or mouse MP121 sequences. The transcripts of MP121 were obtained from liver tissue and code for a protein which shows a considerable amino acid homology to the mature part of the inhibin/activin-like proteins (see Figure 1). The protein sequences of human α -inhibin, inhibin β_A (activin β_A) and inhibin β_B (activin β_B) are described by Mason et al. (Biochem. Biophys. Res. Comm. 135, 957-964 (1986)). Some typical sequence homologies which are specific for known inhibin sequences were also found in the propeptide part of MP121

while other parts of the propeptide of MP121 show considerable differences to inhibin propeptides.

However previous findings show that there are differences between the pattern of expression of MP121 and that of the activins. While activins are mainly expressed in the gonads (activin β_A in ovaries and activin β_B in testes and ovaries), MP121 is mainly expressed in the liver. However up to now the sensitivity of the experiments has not been sufficient to also detect a slight expression. Thus in the case of activins it has for example been described in the literature that expression can also be detected outside the gonads in various rat tissues in adult animals (Meunier et al., Proc. Natl. Acad. Sci. USA 85, 247-251 (1988)) as well as during embryonic development (Roberts et al., Endocrinology 128, 3122-3129 (1991)). Therefore it is also possible that expression of MP121 in other tissues may yet be detected.

Because of the predominant expression of MP121 in liver the expression in one typical cell type of the liver was investigated in more detail. It was shown that the mRNA is expressed abundantly in cultured primary rat hepatocytes as well as in liver cell lines such as HepG2 (ATCC HB 8065). The expression in primary cells is markedly reduced by EGF (Epidermal Growth Factor) treatment after 60 hours. This pattern is completely different compared to activin β_A mRNA, which is barely expressed in hepatocytes but increased drastically after EGF treatment (Yasuda et al., J.Clin.Invest. Vol.92, 1491-1496 (1993)). Likewise, the expression of activin β_A mRNA and MP121 mRNA is reciprocal in remnant rat liver after 70% hepatectomy. MP121 mRNA is detected significantly before hepatectomy but is markedly decreased after 12 hours or later, whereas the mRNA for activin β_A is quite low before but elevated 12 hours or later after hepatectomy. Therefore MP121 seems to have a big influence on the ability of the liver to regenerate and proliferate. The control of MP121 mRNA expression and/or the amount of MP121 protein in liver can be

000107-22218900

In addition the present invention concerns a vector which contains at least one copy of a DNA molecule according to the invention. In such a vector the DNA sequence according to the invention is preferably linked operatively with an expression control sequence. Such vectors are suitable for producing TGF- β -like proteins in stably or transiently-transformed cells. Various animal, plant, fungal and bacterial systems can be used for the transformation and the subsequent culture. The vectors according to the invention preferably contain sequences necessary for replication in the host cell and they are autonomously replicable. In addition the use of vectors is preferred which contain selectable marker genes by which means the transformation of a host cell can be detected.

In addition the invention concerns a protein of the TGF- β family which is coded by a DNA sequence according to claim 1. The protein according to the invention preferably has the amino acid sequence shown in SEQ ID NO. 2 or in SEQ ID NO. 4 or if desired functional parts thereof (as defined above) and exhibits biological properties such as tissue-inductive properties which may be relevant for a therapeutic application. The above-mentioned features of the protein can vary depending on the formation of homodimers or heterodimers with other proteins having the "cystine knot motif" and in particular TGF- β proteins. Such structures may also prove to be suitable for clinical applications and thus are also a subject matter of the present invention. Preferred

heterodimers include heterodimers composed of a monomer of the protein according to the invention and monomers of the α , β_A or β_B inhibin chains. The properties resulting from heterodimer formation can be shifted more towards the properties of activin or inhibins. If for example a heterodimer is formed with inhibin α proteins or with other inhibin β proteins, then it is assumed that the MP121/inhibin (α chain) or MP 121/activin (β_A or β_B chain) heterodimer can inhibit or activate the formation of follicle-stimulating hormone (FSH). MP121/activin heterodimers may also for example influence mesoderm development. Furthermore it is expected that heterodimeric forms with a member of the BMP group of TGF- β proteins lead to an amplification of BMP-like activities such as for example the ability to induce or promote bone formation, formation of cartilage or formation of connective tissue.

The invention therefore also concerns heterodimeric proteins of a protein of the TGF- β family according to the invention which is coded by a DNA sequence as claimed in claim 1 containing a monomer of a protein with the "cystine knot motif" preferably of another member of the TGF- β family. Similar heterodimeric proteins are described in W093/09229, EP 0 626 451 A2 and J. Biol. Chem. 265 (1990), 13198-13205.

In addition the invention concerns chimeric proteins which have functional derivatives or parts of a protein coded by a DNA sequence according to the invention preferably as shown in SEQ ID NO.2 or SEQ ID NO.4, in particular functional parts of the mature protein and additionally parts of another protein. In this case the other protein can also be a protein with a "cystine knot motif" which is preferably also a member of the TGF- β family such as e.g. especially MP52 (PCT/EP94/02630). However, parts of a complete different protein can also be present e.g. receptor-binding domains of proteins which lend the initial MP121 protein another specificity.

Activin A and TGF- β 1, TGF- β 2 and TGF- β 3 have been described to promote survival of dopaminergic neurones in vitro (Krieglstein et al., EMBO J. 14, 736-742 (1995) and Krieglstein et al., Neuroscience 63, 1189-1196 (1994)). In the case of partially purified MP121 it could be shown that the survival of dopaminergic neurones in a 8-day culture is promoted to a greater extent than by the influence of the control supernatant (Figure 5).

During the development of the visual system a projection of axons from the retinal ganglion cells to the special regions in the brain is established. It was shown by several groups that soluble factors isolated from visual areas of the brain can trophically stimulate retinal ganglion cells (Nurcombe, V. & Bennett, M.R., Exp. Brain Res. 44, 249-258 (1981), Hyndman, A.G., Adler, R., Dev. Neurosci. 5, 40-53 (1982), Turner, J.E. et al., Dev. Brain Res. 6, 77-83 (1983), Carri, N.G. & Ebendal, T., Dev. Brain Res. 6, 219-229 (1983)). The formation of nerve fibre fascicles, which most likely represent optic axons stemming from the retinal ganglion cells, depends on neurotrophic factors. Using MP121, a strong stimulation of retinal nerve fibre outgrowth in explant cultures of the embryonic chicken retina was evident as shown in Tab.1 and Figure 7. During these experiments, other members of the TGF- β superfamily, as for example MP52 (DE 195 25 416.3), were also proven to be active.

Results on rat hepatocytes in primary cultures indicate that partial purified MP121 inhibits the initiation of DNA synthesis (Figure 8). The effect of MP121 resembles that of Activin A and TGF- β (Yasuda et al., J. Clin. Invest. Vol. 92, 1491-1496 (1993)) but the concentrations of MP121 which are necessary to block the growth promoting actions of EGF are higher. Nevertheless it can be assumed that MP121 can influence liver growth. Therefore it can be useful in several liver diseases including liver carcinomas.

Activin A is furthermore known for its ability to promote the differentiation of Friend erythroleukemic cells (F5-5) wherefore it was also designated Erythroid differentiation factor (EDF) (Eto et al., Biochem. Biophys. Res. Com. 142, 1095-1103 (1987)). Partial purified MP121 shows also a slight activity in this assay system. Therefore MP121 can be useful in stimulation of erythropoiesis.

The present invention in addition concerns a process for the production of a protein of the TGF- β family which is characterized in that a host cell transformed with a DNA

according to the invention or with a vector according to the invention is cultured and the TGF- β protein is isolated from the cell or/and the culture supernatant. Such a process comprises culturing the transformed host cell in a suitable culture medium and purifying the TGF- β -like protein formed. In this way the process enables the production of an adequate amount of the desired protein for use in medical treatment or in applications using cell culture techniques in which growth factors are needed. The host cell can be a bacterium such as Bacillus or E. coli, a fungi such as yeast, a plant cell such as tobacco, potato or arabidopsis or an animal cell, especially a vertebrate animal cell line such as Mo, Cos or CHO cell lines or an insect cell line. Using the Baculovirus system, expression can also be performed in insect larvae. When producing in bacteria it is possible that the protein according to the invention is produced in the form of inclusion bodies. These inclusion bodies are then renatured according to known methods and the protein is then obtained in an active form (see e.g. Jaenicke, R. and Rudolph, R., Protein Structure, ed. Creighton, T.E., IRL Press, chapter 9). For the production of heterodimeric proteins with other members of the TGF- β family, both protein monomers are expressed either in the same cell or separate in the course of which a common renaturation seems suitable with formation of inclusion bodies. Viral systems such as e.g. the Baculovirus system or the Vaccinia virus system are in particular suitable when coexpressing in the same cell. The production of heterodimeric proteins is in principle known to a person skilled in the art and is described for example in W093/09229 and EP 0 626 451 A2.

The production of chimeric proteins containing other protein parts requires a corresponding change at the DNA level which is familiar to a person skilled in the art and can be carried out by him (EMBO J. 10 (1991), 2105-2110; Cell 69 (1992), 329-341; J. Neurosci. 39 (1994), 195-210).

Yet another subject matter of the present invention is the provision of pharmaceutical compositions which contain a pharmaceutically effective amount of a TGF- β -like protein according to the invention as the active substance. If desired, such a composition comprises a pharmaceutically acceptable carrier or auxiliary substance, diluent or filling agent. Such a pharmaceutical composition can be used alone or in combination with other active substances for example other proteins of the TGF- β family or growth factors such as EGF (epidermal growth factor) or PDGF (platelet derived growth factor) in wound healing and tissue regeneration. Furthermore such a pharmaceutical composition can be used for the prevention of diseases.

Further subject matters are pharmaceutical compositions which contain heterodimeric proteins or/and chimeric proteins according to the invention.

The pharmaceutical composition according to the invention is preferably used for the treatment and prevention of damage to bones, cartilage, liver, connective tissue, skin, mucous membranes, endothelium, epithelium, neurones, kidneys or teeth, for application in dental implants, for application in wound healing or tissue regeneration processes, induction of the proliferation of precursor cells or bone marrow cells, for the maintenance of a state of differentiation and for the treatment of disturbances in fertility or for contraception.

Furthermore the pharmaceutical composition according to the invention can be useful for the treatment of diseases concerning the metabolism, such as digestive disorders or disorders concerning the level of bloodsugar.

A further possible clinical application of the TGF- β -like protein according to the invention is the use as a suppressor of immunoreactions in order to avoid rejection of organ transplants or use in connection with angiogenesis.

Furthermore the protein according to the invention can be used to increase fertility or in contraception. The pharmaceutical composition according to the invention can also be used prophylactically or in cosmetic surgery. Furthermore the application of the composition is not limited to humans but can also include animals in particular pets and domestic animals.

Thus the part of the other protein or other monomer can be used to vary the scope of applications and specificity of heterodimeric proteins and chimeric proteins as desired.

In general diseases which are associated with the expression of MP121 can be treated using the proteins according to the invention either by increasing the amount or activity of MP121 which is present or by suppressing the MP121 activity. Thus the invention also concerns the production of antisense nucleic acids and ribozymes which inhibit the translation of MP121. This inhibition can either be achieved by masking the mRNA with an antisense nucleic acid or by cleavage with a ribozyme.

The production of antisense nucleic acids is known (Weintraub, H.M., Scientific American 262: 40 (1990)). The antisense nucleic acids hybridize with the respective mRNA and form a double-stranded molecule which can then no longer be translated. The use of antisense nucleic acid is for example known from Marcus-Sekura, C.J., Anal. Biochem. 172 (1988), p. 289-295.

Ribozymes are RNA molecules which are able to specifically cleave other single-stranded RNA molecules similar to DNA restriction endonucleases. The production of ribozymes is described in Cech, J. Amer. Med. Assn. 260 (1988), p. 3030.

In this connection it is also possible according to the invention to transfect suitable vectors containing the DNA

MP121 antisense polynucleotides can also be introduced into cells which exhibit an undesired expression of MP121.

Thus within the scope of the invention the receptors for MP121 on cells are also of interest. In order to find receptors, firstly various cell lines can be tested for their binding properties with respect to radioactively labelled MP121 (¹²⁵I-MP121) with subsequent cross-linking. A cDNA library can subsequently be established in an expression vector (obtainable from InVitrogen) from cells which bind MP121. Cells which have been transfected with receptor cDNA can then be selected by the binding of radioactively labelled MP121. These are methods known to a person skilled in the art and have for example been used to isolate activin (Mathews, L.S. & Vale, W.W., Cell 65 (1991), 973-982) and TGF- β type II receptors (Lin, H.Y. et al., Cell 68 (1992), 775-785). In analogy to known activin receptors, the MP121 receptor is also presumably a receptor complex which belongs to this family so that further methods known to a person skilled in the art, such as e.g. PCR with degenerate oligonucleotides, can be used to find parts of the heteromeric complex. This method has also been used for example with the activin and TGF- β type I receptors (Tsuchida et al., Proc. Natl. Acad. Sci. USA 90 (1993), 11242-11246; Attisano et al., Cell 75 (1993), 671-680; Franzén et al., Cell 75 (1993), 681-692).

Finally the present invention concerns an antibody which can bind specifically to the proteins according to the invention or such an antibody fragment (e.g. Fab or Fab'). Processes for

the production of such a specific antibody or antibody fragment are part of the general knowledge of an average person skilled in the art. Such an antibody is preferably a monoclonal antibody. Such antibodies or antibody fragments can also be suitable for diagnostic methods.

In addition it is intended to illustrate the invention by the following examples.

Example 1

Isolation of MP121

1.1 Total RNA was isolated from human liver tissue (40 year old man) according to the method of Chirgwin et al. (Biochemistry, 18, 5294-5299 (1979)). Poly (A+)-RNA was separated from the total RNA by oligo (dT) chromatography according to the manufacturer's instructions (Stratagene poly (A) Quick columns).

1.2 For the reverse transcription reaction 1 to 2.5 µg poly (A+) RNA was heated for 5 minutes to 65°C and quickly cooled on ice. The reaction mixture contained 27 U RNA-Guard (Pharmacia), 2.5 µg oligo (dT)₁₂₋₁₈ (Pharmacia), 5 x buffer (250 mmol/l Tris/HCl pH 8.5, 50 mmol/l MgCl₂, 50 mmol/l DTT, 5 mmol/l of each dNTP, 600 mmol/l KCl) and 20 U AMV reverse transcriptase (Boehringer Mannheim) per µg poly (A+) RNA. The reaction mixture (25 µl) was incubated for 2 hours at 42°C. The cDNA pool was stored at -20°C.

1.3 The deoxynucleotide primers OD and OID shown in Figure 2 were prepared on an automatic DNA synthesizer (Biosearch). Purification was carried out by means of denaturing polyacrylamide gel electrophoresis and isolating the main bands from the gel by isotachopheresis. The oligonucleotides were designed by comparing nucleic acid sequences of known members of the TGF-β family and selecting regions with high conservation. A comparison of this region is shown in Figure

1.4 In the PCR reaction cDNA corresponding to 20 ng poly (A+) RNA were used as starting material (see 1.2) The reaction was carried out in a volume of 50 μ l and contained 1 x PCR buffer (16.6 mmol/l $(\text{NH}_4)_2\text{SO}_4$, 67 mmol/l Tris/HCl pH 8.8, 2 mmol/l MgCl_2 , 6.7 μ mol/l EDTA, 10 mmol/l β -mercaptoethanol, 170 μ g/ml bovine serum albumin (Gibco), 200 μ mol/l of each dNTP (Pharmacia), 30 pmol of each oligonucleotide (OD and OID) and 1.5 U Taq polymerase (AmpliTaq, Perkin Elmer Cetus). The reaction mixture was overlayed with paraffin and 40 PCR cycles were carried out. The products of the PCR reaction were purified by means of phenol/chloroform extraction and concentrated by ethanol precipitation.

1.5 Half of the PCR reaction products was cleaved with the restriction enzymes SphI (Pharmacia) and AlwNI (Biolabs) according to the manufacturer's instructions. The other half was cleaved in a series of reactions using Ava I (BRL), AlwN I (Biolabs) and Tfi I (Biolabs). The restrictions were carried out in 100 μ l using 8 U enzyme for 2 to 12 hours at 37°C (apart from Tfi I at 65°C).

1.6 The products of the restriction cleavage were fractionated by means of agarose gel electrophoresis. After staining with ethidium bromide, uncleaved amplification products were cut out of the gel and isolated by phenol extraction. The DNA obtained was subsequently purified twice by phenol/chloroform extraction.

1.7 A quarter or a fifth of the isolated DNA was reamplified after ethanol precipitation using the same conditions as for the primary amplification except that the number of cycles was reduced to 13. The reamplification products were purified, cleaved with the same enzymes as above and uncleaved products

were isolated from the agarose gels as elucidated above for the amplification products. The reamplification step was repeated.

1.8 After the last isolation from the gel, the amplification products were cleaved by 4 U Eco RI (Pharmacia) under the conditions recommended by the manufacturer. A quarter of the restriction mixture was ligated into the vector pBluescript SK+ (Stratagene) which had been cleaved with Eco RI. After ligation, 24 clones of each enzyme combination were analyzed further by sequencing. There were no new sequences in the mixture which had been cleaved with AlwN I and Sph I, it contained only BMP6 and inhibin BA sequences. 19 identical new sequences, named MP121, were found in the mixtures cleaved with Ava I, AlwN I and Tfi I. These plasmids were named pSK-MP121 (OD/OID). One sequence differed by two nucleotides from this sequence that was otherwise found. Ligation and transformation in *E. coli* was carried out as described in Sambrook et al., Molecular Cloning: A Laboratory Manual (1989).

by 7
The clone was extended to the 3' end of the cDNA according to the method described in detail by Frohmann (published by Perkin-Elmer Corp., Amplifications, 5, 11-15 (1990)). The same liver mRNA which had been used to isolate the first MP121 fragment was reversely transcribed as described above using oligo dT (16mer) linked to the adapter primer (AGAATTCGCATGCCATGGTTCGACGAAGC -T₁₆). The amplification was carried out using the adapter primer (AGAATTCGCATGCCATGGTTCGACG) and an internal primer (GGCTACGCCATGAACTTCTGCATA) prepared from the MP121 sequence. The amplification products were prepared using a further internal primer (ACATAGCAGGCATGCCTGGTATTG) prepared from the MP121 sequence and with the adapter primer. After restriction with Sph I the reamplification products were cloned into the vector pT7/T3 U19 (Pharmacia) which had been cleaved in the same way and sequenced. The clones were characterized by their

sequence overlap with the already known part of the MP121 sequence. One clone, named p121Lt 3' MP13, was used to isolate a Nco I (made blunt using T4 polymerase)/Sph I fragment. This fragment was cloned into one of the above-mentioned pSK-MP121 (OD/OID) vectors whose OD primer sequence was orientated towards the T7 primer of the pSK multiple cloning site. For this the vector was cleaved with SphI and SmaI. The construct was named pMP121DFus6. It comprises the MP121 sequence from position 922 to 1360 as shown in SEQ ID NO. 1.

1.9 A Dde I fragment of pMP121DFus6, which extends from position 931 to 1304 in SEQ ID NO. 1, was used to screen a human liver cDNA library (Clontech, # HL3006b, lot 36223) as described in detail by Ausubel et al., (Current Protocols in Molecular Biology published by Greene Publishing Associates and Wiley-Interscience (1989)). 24 mixed plaques were picked from 8.1×10^6 phages and separated. From this 10 clones which yielded a positive signal using primer LO2 (ACATAGCAGGCATGCCTGCTATTG) and LO11 (CTGCAGCTGTGTTGGCCTTGAGA) from the Dde I fragment were selected and separated. The cDNA was isolated from the phages by means of an EcoRI restriction and cloned into the pBluescript SK vector which had also been cleaved with EcoRI.

Sequencing of one of the resulting plasmids SK121L9.1 showed that the start codon begins at position 128 of SEQ ID NO. 1 since three stop codons are positioned in-frame in front of this start codon at positions 62, 77 and 92. Mature MP121 starts at position 836 of SEQ ID NO. 1 assuming sequence analogy to other TGF- β proteins which corresponds to amino acid 237 in SEQ NO. 2. The stop codon begins at position 1184 of SEQ ID NO. 1.

Plasmid 9K121L9.1 was deposited at the DSM on the 26.04.1994 under the deposit number 9177.

1.10 Isolation of the MP121 cDNA and genomic DNA from the mouse: The sequence information from the human MP121 sequence was used to isolate the MP121 sequence from the mouse. The methods used for this are all known to a person skilled in the art and are described for example in Current Protocols in Molecular Biology (Ausubel et al., Greene Publishing Associates and Wiley-Interscience, Wiley & Sons, 1987-1995) or in Molecular Cloning (Sambrook et al., second edition, Cold Spring Harbour Laboratory Press 1989). Firstly the primers *ACGAATTCGACGAGGCATCGACTGC* and *GCGTCGACTACCATGTCAGGTATGTC* derived from the human MP121 sequence containing additional restriction cleavage sites at the 5' end (EcoR I or Sal I) were synthesized. These primers were used for amplification on genomic mouse DNA. The 0.35 kb fragment which results was subcloned in the Bluescript vector (Stratagene) and used as a radioactive probe. A λ bank with genomic mouse DNA as well as a bank with cDNA was screened according to standard methods. The cDNA was synthesized from RNA, which had been isolated from mouse liver and cloned into λ gt10 using EcoR I/Not I linkers.

MP121 clones were isolated from the genomic as well as from the cDNA bank. A cDNA containing the complete coding sequence was subcloned into the EcoR I cleavage site of the Bluescript vector SK (Stratagene) and the resulting plasmid SKMP121 mouse was deposited on the 10.05.1995 at the DSM (DSM 9964). Complete sequencing resulted in the sequence shown in SEQ ID NO.3. The start codon begins at position 131 in SEQ ID NO.3 and ends at the stop codon starting at position 1187. The protein derived from the sequence is shown in SEQ ID NO.4. Subcloning and analyzing clones containing MP121 from the genomic bank showed that the MP121 sequence contains an intron in the propeptide part of ca. 5.5 kb. This intron is located between positions 446 and 447 in SEQ ID NO.3. The exon/intron junctions are shown in SEQ ID NO.5.

Example 2

Expression of MP121

It is possible to express MP121 in eukaryotic as well as in prokaryotic systems.

Only the mature part of MP121 was used for expression in prokaryotes. After purification the mature MP121 protein expressed in E. coli as a monomer can then be folded back to form a dimer. In order to simplify purification of MP121, an additional 6 histidines can be attached to the N-terminus of the mature protein which facilitate purification of the protein by binding to nickel-chelate columns.

As an example the mature part of human MP121 (amino acid 237 to 352 in SEQ ID NO.2) with an additional 13 amino acids, including 6 histidines at the N-terminus, (MHIIHHHKKLEFAM) was expressed in the prokaryotic vector pBP4. This vector is a pBR322 derivative having tetracyclin resistance which in addition contains the T7 promoter from the pBluescript II SK plasmid (Stratagene). Furthermore the vector contains a ribosomal binding site following the T7 promoter and a start codon followed by 6 codons for histidine. A terminator (T0) follows after several single restriction cleavage sites such as Eco RI, Xho I, Sma I and Apa I for the insertion of inserts as well as stop codons in all three reading frames. In order to obtain the cDNA for the mature part of MP121, PCR was carried out on the plasmid SK121L9.1 (DSM depositary number: 9177) using the two oligonucleotides GAATTCGCCATGGGCATCGACTGCCAAGGAGG and CCGCTCGAGAAGCTTCAACTGCACCCACAGGC. Both oligonucleotides contain additional restriction cleavage sites at their ends (Eco RI and Nco I or Xho I and Hind III). In an intermediate step the resulting 377 bp fragment was cloned with blunt ends into the pBluescript II SK vector (Stratagene) that had been cleaved with Eco RV. One clone in the orientation of the 5' end of MP121 towards the T7 promoter was cleaved with Eco RI

000000-000000

Sub
D10

and the resulting insert (0.38 kb) was cloned into the pBP4 vector that had also been cleaved with Eco RI. The correct orientation of the insert in the resulting plasmid pBP4MP121His was established by restriction analysis and sequencing. The plasmid pBP4MP121His was deposited on the 30.1.1995 at the DSM (depository number: 9704). The expression of MP121 protein can be achieved by simultaneously providing T7 RNA polymerase. T7 RNA polymerase can be provided by various methods such as e.g. by a second plasmid with a gene for T7 RNA polymerase or by infection with phages which code for T7 RNA polymerase or also by special bacterial strains which have integrated the gene for T7 RNA polymerase. The mature MP121 protein with a His-tag (MP121His) is produced in inclusion bodies by using the bacterial strain BL21 (DE3)pLysS (Novagen, #69451-1) and inducing the T7 RNA polymerase expression with IPTG according to the manufacturer's instructions. In SDS polyacrylamide gels (15 %) the protein exhibits an apparent molecular weight of nearly 16 kD (theoretical molecular weight: 14.2 kD) as is shown representatively in the Western blot of Fig. 3. The bacteria transformed with pBP4 as controls do not show any staining of specific bands. Due to the His-tag this protein can be purified on nickel-chelating agent columns as described for example by Hochuli et al., (BIO/Technology Vol. 6, 1321-1325 (1988)). An additional purification is possible by means of reversed phase HPLC. A reversed phase column (Nucleosil 300-7C4 from Macherey-Nagel, Type 715023) was used with a flow-rate of 2 ml/min and an acetonitrile gradient in 0.1 % TFA of 0 to 90 % within 100 minutes. MP121His elutes under these conditions after ca. 40 % acetonitrile.

The mature part of MP121 (amino acid 237 to 352 in SEQ ID NO.2) was additionally expressed in E.coli with one additional methionine at the N-terminus only using again a system with the T7 RNA Polymerase. The expression level was improved by including a gene for the lacI repressor in the expression plasmid (as it is used in the pET vectors from Novagen) and

using another E. coli strain, HMS 174 (DE3) (Novagen #69453). Inclusion bodies can be isolated by standard methods and washed with 2 M guanidinium chloride /HCL in 20 mM Tris pH 8.0. MP121 is further purified by a reversed phase HPLC as described for MP121His.

Additionally monoclonal antibodies were developed in mice. A peptide of 26 amino acids from the mature part of MP121 was used as an antigen: PLSLLYYDRDSNIVKTDIPDMVVEAC. The antigen was coupled to ovalbumin using the free SH group of the cysteine according to conventional methods. Other constructs could be used as antigens also, as for example the dimeric mature MP121.

Immunization of BALB/c mice was performed according to conventional methods. The coupled peptide was used for example as antigen in combination with complete Freund's adjuvant for the first immunization and in combination with incomplete Freund's adjuvant in successive immunizations. The antigen (5-10 µg each time) was injected subcutaneously in the hind limbs of three mice at day 17, 14, 10, 7, 4 and 1 before the isolation of popliteal lymphatic nodes underneath the knee joint. A suspension of cells was produced for fusion to myeloma cells (P3X63Ag8.653, ATCC, CRL 1580) by the help of PEG. These techniques are described in more detail by Peters, J.H. & Baumgarten, H. (1990, Monoklonale Antikörper - Herstellung und Charakterisierung, Springer Verlag, 2.Auflage). It is possible to select for fused hybridomas by addition of azaserine and hypoxanthine. The supernatants of different wells were tested after 8-10 days with ELISA and Western blot analyses using MP121 expressed in eukaryotic and prokaryotic cells. The cells with the best positive results were further subcloned to isolate cells producing only one monoclonal antibody. To purify the monoclonal antibodies 1 liter of cell culture supernatant containing the monoclonal antibody was produced using roller bottles (Schott) in a "Cell-Roll" (Former Scientific) according to standard methods.

In each case the determination whether it is MP121 protein was carried out by means of Western blot analysis using MP121-specific antibodies. Polyclonal antibodies against MP121 were produced in chicken as well as in rabbits. In order to obtain the antigen for the immunization, a part of the mature part of MP121 (amino acid 260 to 352 in SEQ ID NO.2) was fused with the first 98 amino acids of the polymerase of the MS2 bacteriophage and expressed in E. coli. After isolating the inclusion bodies, the fusion protein (MS2-MP121) was separated on polyacrylamide gels and isolated for the immunization after staining with copper by means of electro-elution (Tessmer, U. & Dernick, R., IBL (1990) 8-13). It is possible to specifically detect the expression of MP121 using antibodies from chicken as well as from rabbits. Chicken antibodies were used for the schematic Western blot in Figure 3 which had been purified further by means of PEG precipitation (Thalley B.S. and Carroll, S.B., BIO/Technology Vol. 8, 934-938 (1990)) and by means of membrane-bound antigen (fusion protein (MS2-MP121)) (18.17 in Sambrook et al., Molecular Cloning, second edition, Cold Spring Harbor Laboratory Press 1989). Anti-chicken IgG coupled to alkaline phosphatase (Sigma A9171) was used as the second antibody. The detection was carried out according to the manufacturer's instructions using the Tropix Western-Light Protein Detection Kit (Serva #WL10RC).

In order to obtain biologically active material, the purified monomeric MP121 expressed in *E. coli* can be folded back to form a dimeric MP121. This can be carried out according to the methods for example described by Jaenicke, R. & Rudolph, R.

(Protein structure, ed. Creighton, T.E., IRL Press, chapter 9).

The Vaccinia virus expression system was used for expression in eukaryotic cells as it is described in detail and in a form which can easily be repeated by a person skilled in the art in Current Protocols in Molecular Biology (Ausubel et al., Greene Publishing Associates and Wiley-Interscience, Wiley & Sons) abbreviated in the following as CP, in chapter 16 unit 16.15-16.18. The system is based on the fact that foreign DNA can be integrated by homologous recombination into the genome of the Vaccinia virus using certain vectors. For this purpose the vector used contains the TK (thymidine kinase) gene from the Vaccinia genome. In order to enable selection for recombinant viruses, the vector additionally contains the E. coli xanthine-guanine-phosphoribosyl transferase gene (gpt) (Falkner, F.G. & Moss, B., J. of Virol. 62 (1988), 1849-1854). The cDNA with the complete region coding for MP121 was cloned into this vector.

PCR reactions and intermediate cloning was necessary in order to shorten the 5' and 3' untranslated regions of the initial plasmid SK121L9.1 (DSM, depositary number: 9177) and to insert single restriction cleavage sites at the ends. All PCR reactions were carried out using the plasmid SK121L9.1 (DSM depositary number: 9177). In order to shorten the 5' untranslated end, the primer CCCGGATCCGCTAGCACCATGACCTCCTCATTGCTTCTG with an inserted Bam HI and NheI restriction cleavage site was used in a PCR with an internal primer (CCCTGTTGTCCCTCTAGAAGTG). In an intermediate step the fragment obtained was cloned into Bluescript SK (Stratagene), sequenced and checked for concurrence with the sequence shown in SEQ ID NO.1. The Sph I/Eco RI fragment (0.22 kb) from the plasmid pBP4MP121His was used to shorten the 3' untranslated end.

Both end fragments of MP121 were linked to the missing middle DNA sequence from the plasmid SK121L9.1 (DSM depositary

number: 9177) by means of internal restriction cleavage sites (Xba I and Sph I) according to standard methods (Sambrook et al. Molecular Cloning, second edition, Cold Spring Harbor Laboratory Press 1989). The shortened cDNA obtained in this way having the complete reading frame for MP121 (nucleotide 128 to nucleotide 1184 in SEQ ID NO.1) could be cloned into the vector pBP1 which had also been cleaved by using the restriction cuts Bam HI and Eco RI. The resulting plasmid pBP1MP121 was deposited on 12.1.95 at the DSM (depository number: 9665).

The plasmid pBP1MP121 was used for the production of recombinant Vaccinia viruses. For this 143B cells (HuTk-, ATCC CRL 8303) which were 80 % confluent were infected with Vaccinia wild-type virus (1 virus per 10 cells) in 1 ml PBS in 35 mm culture plates for 30 minutes at room temperature while shaking occasionally. After aspirating the supernatant and adding 2 ml culture medium (MEM, Gibco BRL #041-01095 containing 1:500 diluted penicillin and streptomycin Gibco BRL #043-05140), they were incubated for 2 hours at 37°C. Subsequently the medium was removed and these cells were transformed for ca. 15 hours at 37°C using 100 ng pBP1MP121, 2 µg carrier DNA (calf thymus, treated with ultrasound, Boehringer Mannheim #104175) and 10 µl Lipofectin (Gibco BRL #18292-011) in 1 ml MEM. After addition of 1 ml MEM containing 20 % FCS (Sigma #F-7524) they were incubated for a further 24 hours at 37°C and subsequently the lysed cells were frozen.

Gpt selection for xanthine-guanine-phosphoribosyl transferase and isolation and amplification of individual recombinant viruses was essentially carried out as described in unit 16.17 of CP with the difference that RK13 cells (ATCC CCL 37) were used.

Integration of the MP121 cDNA into the viral genome was confirmed by dot blot analysis (CP unit 16.18). A recombinant virus from the transfection with pBPMP121 and the wild-type

virus were used for expression analyses in cell lines 143B (HuTk-, ATCC CRL 8303, human) and NIH-3T3 (DSM ACC 59, Swiss mouse embryo). The cells were cultured according to the distributor's instructions. Confluent cells were infected for 30 minutes at 37°C with the three-fold number of viruses and subsequently the respective culture medium containing 10 % FCS and penicillin/streptomycin (1:500, Gibco BRL #043-05140) was added. The medium was removed after 6 hours at 37°C, the cells were washed twice with e.g. HBSS (Gibco BRL #14180-046) and production medium (MEM for HuTk- or DMEM containing 4.5 g/l glucose and NEAA (Gibco BRL #11140-035) for NIH-3T3 each of which contained aprotinin (Fluka #10820, 50 U/ml) and penicillin/streptomycin) without FCS. After a production period of 20 to 22 hours, the cell supernatant was collected. The expression was analysed by means of Western blots according to standard methods (CP unit 10.8). For this the proteins from 1 to 3 ml cell culture supernatant were precipitated by addition of an equivalent volume of acetone and incubating for at least one hour on ice and centrifuged. After resuspending the pellets in application buffer (7 M urea, 1 % SDS, 7 mM sodium dihydrogen phosphate, 0.01 % bromophenol blue and 1 % β -mercaptoethanol if desired) they were separated in 15 % polyacrylamide gels. A pre-stained protein molecular weight standard (Gibco BRL #6041-020) was used as marker proteins. Transfer onto a PVDF membrane (Immobilon #IPVH00010) and blocking the membrane was carried out according to standard methods.

A representative schematic diagram of the results of the Western blot in Figure 3 shows that MP121-specific bands occur in the recombinant virus infected cells. The expression of MP121 in NIH-3T3 cells leads to a secreted protein with an apparent molecular weight in the gel of about 18 kD under non-reducing conditions (expected theoretical molecular weight: 25 kD). Under reducing conditions the protein migrates at about 15 kD in the gel (expected theoretical molecular weight: 12.5 kD). These results show that MP121 is expressed

as a dimeric mature protein as expected. The migration behaviour of the dimeric MP121 protein which is only slightly slower than the monomeric MP121 protein is probably due to its globular structure. The processing of the precursor protein to form the mature protein could also be demonstrated in HuTk-cells. No bands occurred in the Western blot with cells (HuTk- or NIH-3T3) infected with wild-type viruses (without integrated foreign DNA).

Further expression studies of MP121 using the Vaccinia virus system revealed that several cell lines express in addition to the dimeric MP121 also significant amounts of a monomeric form. This monomeric form seems to be folded and has a more globular structure because it runs faster in PAGE/Western blot analyses than the reduced monomer derived from the dimeric MP121 after treatment with DTT. Figure 6 shows the expression of dimeric and monomeric MP121 in HepG2 cells (Hepatocellular carcinoma, human, ATCC HB 8065). A residual unprocessed precursor form appears in addition. It was already shown by our Northern blot analysis that the HepG2 cells naturally transcribe the MP121 gene, therefore it can be assumed that the appearance of monomeric MP121 is of physiological relevance.

The monomeric MP121 was found besides the dimeric MP121 in significant amounts in Mv1Lu (NBL-7, lung, mink, ATCC CCL64) and Hela (Epitheloid carcinoma, cervix, human, ATCC CCL2) too. In addition, MP121 was expressed using the Baculovirus expression system (Invitrogen). After infection of insect larvae (*Trichoplusia ni*) with recombinant viruses, MP121 was detected in the haemolymph after 3-4 days in the dimeric form.

When co-transfection with recombinant Vaccinia viruses that code for various members of the TGF- β family has also taken place, the Vaccinia virus expression system is also particularly suitable for the production of heterodimers. It is then possible to separate heterodimers from homodimers by affinity columns using specific antibodies against the

Sub
D13

Example 3
Investigation of the expression of MP121 in various
mouse tissues

GGATCCGAATTTCGGCTTGGAGTGTATGGCAAGG
and GGATCCGAATTCTTCTGGGACCTGGCAACTCTAG.

Sub
D14

and GCAAGCTTT (GA) TA (TC) TC (GA) TC (GA) TC.

The resulting PCR fragments were subcloned into the vector pGEM-4 (Promega) and tested. The activin-specific and thus in the RPA protected sequences have a fragment size of 369 bp in the case of activin β_A and 254 bp in the case of activin β_B . In MP121 the protected fragment comprises the sequence from position 887 to position 1164 in SEQ ID NO.3. The fragments cloned into pGEM-4 were transcribed in vitro in order to produce radioactively labelled antisense RNA probes. This was

carried out according to the manufacturer's instructions (Promega, Riboprobe Gemini Systems) using 100 μ M CTP and at the same time α^{32} P-CTP (800 Ci/mmol, Amersham). A radioactively labelled RNA was also synthesized as a control from the plasmid pTri-GAPDH (Ambion #7431) linearized with Dde I but using 1 mM CTP. After isolating the 4 antisense RNA probes from polyacrylamide gels, these were incubated at 42°C overnight in the same mixture with the respective tissue RNA from the mouse (10 μ g total RNA per probe having 1×10^5 cpm). It was analyzed in a denaturing gel according to standard methods with a subsequent autoradiography for 4 days.

The analysis of MP121 mRNA expression in liver cells or remnant liver was performed likewise or using Northern blot analysis according to standard procedures (see CP, Chapter 4 or Molecular Cloning, Sambrook et al., 2nd Edition, Cold Spring Harbor Laboratory Press 1989).

Hepatocytes were isolated from rat (Wistar) liver and cultured according to Yasuda et al. (J. Clin. Invest. Vol. 92, 1491-1496 (1993)). The cells were washed prior to incubation with fresh serum-free medium containing 0.1 nM insulin, 0.1% BSA, optionally 1 nM EGF and (1 nM) partially purified MP121 (see Example 4).

Partial hepatectomy (about 70% of the rat liver) was performed as described by Higgins & Anderson (Arch. Pathol. 12, 186-202 (1931)) under ether anesthesia.

Example 4

Partial purification of MP121 and examination of the activity of partially purified MP121

The MP121 protein which had been obtained by expression in the Vaccinia virus system (see example 2) could be partially purified by means of two columns.

In order to produce MP121 confluent NIH-3T3 cells (DSM ACC 59, Swiss mouse embryo) were infected with the same number of

recombinant viruses for 30 minutes at 37°C and subsequently the appropriate culture medium containing 10 % FCS and penicillin/streptomycin was added.

After 4 hours at 37°C the medium was removed, the cells were washed twice and production medium (see Example 2) without FCS was added. After 20 to 22 hours production, the cell supernatant was collected and centrifuged in order to remove the viruses (40000 x g for 30 minutes at 4°C) and filtered (0.1 µm pore size, Millex VV, Millipore # SLVV25LS). The control supernatant (wt) was obtained in a comparable manner after infection by wild-type Vaccinia viruses. The expression of MP121 was checked by means of Western blot analysis and estimated to be 50-100 µg/l.

The cell culture supernatant containing MP121 (1.1 l) was admixed with the protease inhibitor PMSF (1 µM), brought to a final concentration of 1 M $(\text{NH}_4)_2\text{SO}_4$, 20 mM Tris pH 8.0 and loaded onto a phenyl-Sepharose (fast flow (high sub) Pharmacia #17-0973-05) column (5 ml bed) equilibrated in buffer A (1 M $(\text{NH}_4)_2\text{SO}_4$, 20 mM Tris pH 8.0). The loaded column was washed with 15 column volumes of buffer A and 10 column volumes of buffer B (20 mM Tris pH 8.0) and eluted within 50 minutes (5 ml per fraction) with a linear gradient to 100 % buffer C (20 mM Tris pH 8.0, 80 % ethylene glycol) at a flow rate of 1 ml/min. It was possible to check that MP121 eluted between 50 and 80 % ethylene glycol by means of Western blot analysis. Aliquots of these fractions were examined using 15 % polyacrylamide silver-stained gels according to the manufacturer's instructions (Silver Stain-II, Daiichi #SE140000) and the fractions containing MP121 were pooled. After purification of the control supernatant comparable fractions were also pooled after analysis in silver-stained gels.

The pooled fractions were purified further with the aid of reversed phase HPLC. For this a C8 column (Aquapore RP300, Applied Biosystems, particle size: 7 µm, pore size: 300Å) was

equilibrated with buffer A (0.1 % trifluoroacetic acid/water). After loading the column with the pooled fractions containing MP121 from the phenyl-Sepharose column, it was extensively washed with buffer A. The bound protein was eluted at a flow rate of 0.2 ml/min using a linear gradient of 1.5 % buffer B (90 % acetonitrile, 0.1 % trifluoroacetic acid) per minute. Fractions of 600 μ l were collected and analyzed in a Western blot as well as with silver-stained gels. Under the selected conditions MP121 protein eluted after about 55% acetonitrile. The fractions containing MP121 were pooled. The same was carried out with the corresponding fractions from the purification of the control supernatant. The analysis in the silver gel showed that MP121 was still contaminated by other proteins. Further purification steps are necessary to obtain pure MP121.

Other methods known to a person skilled in the art such as gel sieve columns, ion exchange columns, affinity columns or metal chelate columns could also be used for the further purification.

It was estimated from Western blot analysis that ca. 8 μ g partially purified MP121 was obtained from 1 l of cell culture supernatant. The partially purified protein was stored lyophilized at -80°C.

In order to investigate the influence of MP121 on dopaminergic neurones, neurones from the mesencephalic floor of 14 day-old rat embryos (E14) were isolated according to a method described by Shimoda et al. (Brain Res. 586, 319-331 (1992)). The cells were singled out and cultured as described by Kriegstein et al., (Neuroscience 63, 1189-1196 (1994)). The cell density on polyornithine/laminin-coated cover glasses is 200000 cells/cm². After culture for 24 hours and subsequently every three days two-thirds of the medium (500 μ l) was removed and replaced by fresh medium containing the respective additives. The lyophilized MP121 partially purified by phenyl

sepharose and reversed phase HPLC was dissolved in 50 % acetonitrile and added to the medium. The final concentration of MP121 in the medium is 20 ng/ml (the final concentration of acetonitrile is 0.3 %). A comparable amount from the control supernatant (wt) which had been purified in a comparable manner was dissolved in 50 % acetonitrile and added. The medium control also contains 0.3 % acetonitrile. After eight days the cultures were fixed for 10 minutes at room temperature in 4 % paraformaldehyde; the cells were made permeable with acetone (10 min, -20°C) and washed with PBS (phosphate buffered saline). After treatment with 1 % H₂O₂ in PBS, washing and blocking with horse serum, they were stained immunocytochemically. Tyrosine hydroxylase (TH) is a limiting enzyme in the biosynthesis of dopamine and other catecholamines so that TH can be used as a marker for dopaminergic neurones in the present cultures (cells containing noradrenaline were not isolated). TH was detected by a 1 hour incubation at 37°C using a mouse-monoclonal antibody against rat TH (diluted 1:200, Boehringer Mannheim) and subsequent detection using the Vectastair ABC kit (Vecto Labs). TH-positive cells were counted in an area of 0.12 cm². It can be seen from Fig. 5 that MP121 has a positive effect on the survival of dopaminergic neurones.

In order to investigate the neural influence of MP121 in another system explant cultures of the embryonic retina were used. This organotypic culture system is described in detail by Carri, N.G. & Ebendal, T. (Dev. Brain Res. 6, 219-229 (1983)), Carri, N.G. & Ebendal, T. (Anat. Rec. 214, 226-229 (1986) and Carri, N.G. et al. (J. Neurosci. Res. 19, 428-439 (1988))). This assay measures the stimulation of extending nerve fibres from the embryonic retina on a collagen substratum. Briefly, the retinal explants were taken from the chick retina (White Leghorn, embryonic day 6) and the neural retina was separated from the pigment epithelium and mesenchymal cells by repeated washing. The organotypic explants were transferred to collagen-coated culture dishes

and incubated overnight (37.5°C, 5% CO₂). The lyophilized MP121 partially purified by Phenyl-Sepharose and reversed phase HPLC was dissolved in aqueous buffer or 50% acetonitrile and diluted in the culture medium to a final concentration of 1.25 ng/ml, 12.5 ng/ml, 25 ng/ml, 50 ng/ml, 100 ng/ml, 200 ng/ml, whereby it makes no difference in the results whether acetonitrile or aqueous buffers were used for solubilization. A comparable amount from the control supernatant (wt) which had been purified in a comparable manner was added in control assays. For the background fibre outgrowth, standard tissue culture medium with only bovine serum added was used. The incubation was continued and after a 4 day period in culture the maximum length of the leading fascicles was measured in an inverted microscope under dark-field illumination. As shown in Table 1, MP121 dose-dependently stimulated the outgrowth of nerve fibres being maximally active at about 25 ng/ml resulting in a real fibre length of about 1.7 mm. Figure 7 shows the fibre outgrowth in a living culture after treatment with MP121 (5 ng/ml). The control (wt) did not stimulate fibre outgrowth as tested in concentrations equivalent to those used for the active MP121.

MP121 (ng/ml)	Length (units)	Mean±SEM
1.25	7/12/5/6	7.5±1.5
12.5	19/20/13/26	19.5±2.6
25	50/52/60/71/65/53	58.5±3.4
50	37/32/48/41/36/20	35.6±3.8
100	21/8/19/18	16.5±2.9
200	11/8/12/10	10.2±0.8

Table 1: Retinal neurite length after 4 days in culture treated with different concentrations of MP121. The neurite lengths of the background fibre outgrowth in the control tissue culture medium were 5.5/8/10/11/4.8/7 units giving a mean of 7.7 units (SEM 1.00). The neurite lengths of the wt control

(used in equivalent concentrations as MP121) was in the same range as the background fibres. Each unit represents 0.03 mm real scale in the culture dish.

In order to investigate the influence of MP121 on liver derived cells hepatocytes were isolated from rat (Wistar) liver and cultured according to Yasuda et al. (J. Clin. Invest. Vol. 92, 1491-1496 (1993)). The cells were washed prior to incubation with fresh serum-free medium containing 0.1 nM insulin, 0.1% BSA and 1 nM EGF. The lyophilized MP121 partially purified by phenyl sepharose and reversed phase HPLC was solubilized in acetonitrile as usually and added to the medium at various concentrations (see Figure 8). A comparable amount from the control supernatant (wt) which had been purified in a comparable manner was used as a control. The hepatocytes were incubated for 72h and 0.5 μ Ci [3 H]Thymidine/ml was included for the last 24 hours as described by Mead & Fausto (Proc.Natl.Acad.Sci.USA 86, 1558-1562 (1989)). [3 H]Thymidine incorporation into trichloroacetic acid-precipitable material was subsequently measured as described by McNeil et al. (J.Cell Biol. 101, 372-379 (1985)).

In order to investigate the influence of MP121 on erythroid differentiation its influence on Friend leukemia cells (F5-5) was measured. Therefore Friend leukemia cells were cultured in microtiter plates essentially as described by Eto et al. (Biochem. Biophys. Res. Com. 142, 1095-1103 (1987)). The lyophilized MP121 partially purified by phenyl sepharose and reversed phase HPLC was solubilized as already described, added to the Friend cells at various concentrations (see Figure 9) and incubated for 5 days. The percentage of differentiated cells was determined after staining with o-dianicidine.